

Clock model interpolation and symmetry breaking in $O(2)$ models¹

Leon Hostetler¹, Jin Zhang², Ryo Sakai²,
Judah Unmuth-Yockey³, Alexei Bazavov¹, and Yannick Meurice²

¹ Michigan State University

²University of Iowa

³Fermilab

May 10, 2021
QuLAT Collaboration

¹On arXiv soon!

Outline

- 1 Motivation
- 2 The Models
 - Extended- $O(2)$ model
 - Extended- q clock model
- 3 MC results from the extended- q clock model
- 4 TRG results from the extended- q clock model
- 5 Characterizing the phase transitions
 - \rightarrow over to Jin

Motivation

- Ultimately, we want to do quantum simulation of lattice QCD
- First, we must be able to do this with simpler Abelian models
- Given the limited number of qubits available, it is important to optimize the discretization procedure
- One can make a \mathbb{Z}_q approximation of a continuous $U(1)$ symmetry
- To optimize such a \mathbb{Z}_q approximation, it is useful to build a continuous family of models that interpolate among the various possibilities
- This brings us to our extended- $O(2)$ model

The Extended-O(2) Model

- We consider an extended-O(2) model in 2D with action

$$S_{\text{ext-O}(2)} = \sum_{x,\mu} \cos(\varphi_{x+\hat{\mu}} - \varphi_x) + \gamma \sum_x \cos(q\varphi_x)$$

- When $\gamma = 0$, this is the classic XY model, with a BKT transition
- When $\gamma > 0$, the second term breaks periodicity and we must choose $\varphi \in [\varphi_0, \varphi_0 + 2\pi)$ for some choice φ_0
- When $\gamma \rightarrow \infty$, the continuous angle φ is forced into the discrete values

$$\varphi_0 \leq \varphi_{x,k} = \frac{2\pi k}{q} < \varphi_0 + 2\pi$$

- ▶ For $q \in \mathbb{Z}$, this is the ordinary q -state clock model with \mathbb{Z}_q symmetry
- ▶ For $q \notin \mathbb{Z}$, this defines an interpolation of the clock model for noninteger q

The Extended- q Clock Model


- In the limit $\gamma \rightarrow \infty$, we can replace the action with

$$S_{\text{ext-}q} = \sum_{x,\mu} \cos(\varphi_{x+\hat{\mu}} - \varphi_x)$$

- We directly restrict the previously continuous angles to the discrete values

$$\varphi_0 \leq \varphi_{x,k} = \frac{2\pi k}{q} < \varphi_0 + 2\pi$$

- We choose $\varphi_0 = 0$, i.e. $\varphi \in [0, 2\pi)$, but we also investigate $\varphi_0 = -\pi$
- For $q \notin \mathbb{Z}$, divergence from ordinary clock model behavior is driven by the introduction of a “small angle”:



Monte Carlo results at small volume




Figure: Results for the extended- q clock model from a heatbath algorithm on a 4×4 lattice.

Phase transitions for $q \in \mathbb{Z}$




Figure: For integer $q \geq 5$, both peaks in the specific heat are associated with BKT transitions (see Li et. al. in Phys. Rev. E **101**, 060105(R) (2020)).

Phase transitions for $q \notin \mathbb{Z}$ (?)




Figure: For non-integer q , are these also BKT transitions or are they something else?

TRG

- In the Monte Carlo approach, we use a Markov chain importance-sampling algorithm to generate equilibrium configurations
 - ▶ Unfortunately, in this model for non-integer q and large β , the configuration space splits into two sectors and the Markov chain tends to get stuck in one or the other
 - ▶ Monte Carlo has difficulty sampling this model appropriately at $\beta > 1$ for $q \notin \mathbb{Z}$
 - ▶ Integrated autocorrelation time explodes, and we have to perform billions of heatbath sweeps already on a 4×4 lattice
 - ▶ Studying this model on larger lattices with Monte Carlo is challenging
- Tensor renormalization group (TRG) approach can be used instead
 - ▶ We validate TRG against Monte Carlo in the regime accessible to Monte Carlo
 - ▶ Then we use TRG to explore lattice sizes and β -values beyond the reach of Monte Carlo

TRG results at large volume




Figure: Specific heat results for the extended- q clock model from TRG obtained by Ryo for $q = 4.1, 4.5, 4.9,$ and 5.0 at volumes from $2^2 \times 2^2$ up to $2^{10} \times 2^{10}$.

Characterizing the phase transitions

→ now over to Jin for his results on the phase transitions

Thank you!

Additional Slides:

Overview

- 6 The Extended-O(2) Model
 - Monte Carlo results
- 7 The Extended- q Clock Model
 - Definitions
 - TRG Energy Density
 - Choice of φ_0
 - Effect of Angle Cutoff
 - TRG Validation
 - Autocorrelation
 - Monte Carlo results
- 8 Phase Diagrams

Some Monte Carlo Results




Figure: Results from a Metropolis algorithm on a 4×4 lattice with $q = 5.0$.

Some Monte Carlo Results




Figure: Results from a Metropolis algorithm on a 4×4 lattice with $\gamma/\beta = 5.0$.

Overview

- 6 The Extended-O(2) Model
 - Monte Carlo results
- 7 The Extended- q Clock Model
 - Definitions
 - TRG Energy Density
 - Choice of φ_0
 - Effect of Angle Cutoff
 - TRG Validation
 - Autocorrelation
 - Monte Carlo results
- 8 Phase Diagrams

Definitions

Internal energy

$$\langle E \rangle = -\frac{\partial}{\partial \beta} \ln Z = -\langle S \rangle$$

Specific heat

$$C = \frac{1}{V} \frac{\partial \langle E \rangle}{\partial T} = \frac{\beta^2}{V} (\langle S^2 \rangle - \langle S \rangle^2)$$

Magnetization

$$\langle \vec{M} \rangle = \frac{1}{\beta} \frac{\partial}{\partial \vec{H}} \ln Z = \left\langle \sum_x \vec{s}_x \right\rangle$$

Magnetic susceptibility

$$\chi_{\vec{M}} = \frac{1}{V} \frac{\partial \langle \vec{M} \rangle}{\partial \vec{H}} = \frac{\beta}{V} \left(\langle |\vec{M}|^2 \rangle - |\langle \vec{M} \rangle|^2 \right).$$

Proxy magnetization and susceptibility

$$\langle |\vec{M}| \rangle = \left\langle \left| \sum_x \vec{s}_x \right| \right\rangle, \quad \chi_{|\vec{M}|} = \frac{\beta}{V} \left(\langle |\vec{M}|^2 \rangle - \langle |\vec{M}| \rangle^2 \right)$$

Energy Density from TRG





Figure: Log of the energy density from TRG with volume 1024×1024 .

Choice of φ_0

- Choice of φ_0 can change the DOF in the model
- We choose $\varphi_0 = 0$, i.e. $\varphi \in [0, 2\pi)$, but we also investigate $\varphi_0 = -\pi$



Effect of Angle Cutoff




Figure: Specific heat for $q = 3.141592654$ with different cutoffs for the allowed angles. When the cutoff is removed (i.e. $\varphi \in (-\infty, \infty)$), all irrational q reduce to the XY model, and all rational $q = r/s$ reduce to the r -state ordinary clock model.

TRG Validation




Figure: Comparison of specific heat calculated via exact and TRG for $q = 4.0$. Bottom panels show difference of TRG from exact.




Figure: Comparison of specific heat calculated via MC and TRG for $q = 4.9$. Bottom panels show difference of TRG from MC with MC as baseline.

Autocorrelation

$$\tilde{\tau}_{X,int} = 1 + 2 \sum_{t=1}^T \frac{C(t)}{C(0)} \quad \text{where} \quad C(t) = \langle X_i X_{i+t} \rangle - \langle X_i \rangle \langle X_{i+t} \rangle$$




Figure: Integrated autocorrelation time for the energy density on a 4×4 lattice using heatbath algorithm.




Figure: The values of T used to extract τ_{int} .

Some Monte Carlo results for $\gamma \rightarrow \infty$




Figure: Results from a heatbath algorithm on a 4×4 lattice.

Some Monte Carlo results for $\gamma \rightarrow \infty$




Figure: Results from a heatbath algorithm on a 4×4 lattice.

Some Monte Carlo results for $\gamma \rightarrow \infty$




Figure: Results from a heatbath algorithm on a 4×4 lattice.

Some Monte Carlo results for $\gamma \rightarrow \infty$




Figure: Results from a heatbath algorithm on a 4×4 lattice.

Some Monte Carlo results for $\gamma \rightarrow \infty$





Figure: Results from a heatbath algorithm on a 4×4 lattice.


Overview

- 6 The Extended-O(2) Model
 - Monte Carlo results
- 7 The Extended- q Clock Model
 - Definitions
 - TRG Energy Density
 - Choice of φ_0
 - Effect of Angle Cutoff
 - TRG Validation
 - Autocorrelation
 - Monte Carlo results
- 8 Phase Diagrams

Phase diagram for $\gamma = \infty$ and $\varphi_0 = 0$



Phase diagram for $\gamma = \infty$ and $\varphi_0 = -\pi$



Phase diagram for finite γ

